

Illinois Department of Natural Resources
Division of Fisheries

Nearshore Fish Community of Lake Michigan: 2014 Summer Assessment

Steven R. Robillard

Illinois Department of Natural Resources

Lake Michigan Program

March 13, 2015

This work was funded by the Federal Aid in Sport Fish Restoration program (F-65-R)

Table of Contents

Executive Summary.....	1
Introduction.....	2
Methods.....	3
Results and Discussion.....	4
Acknowledgements.....	7
Literature Cited.....	8
Tables.....	9-13
Figures.....	14-16

List of Tables

Table 1. Amount of electrofishing effort (hrs:min) and water temperatures in four Illinois harbor areas sampled in 2014.

Table 2. Proposed lengths (mm) for smallmouth and largemouth bass of various size categories (from Gabelhouse 1983).

Table 3. Fish species sampled during summer 2014 by electrofishing in three Illinois harbors and along the shoreline in Calumet Harbor. A = abundant or frequently sampled; P = few specimens observed or infrequently sampled.

Table 4. Smallmouth bass stocking in Lake Michigan harbors, 1969–1985.

Table 5. Catch-per-unit-effort (No. fish / 1 hr electrofishing) of Stock-size (≥ 180 mm) smallmouth bass in four Illinois harbors, 1999-2014.

Table 6. Catch-per-unit-effort (No. fish / 1 hr electrofishing) of Stock-size (≥ 200 mm) largemouth bass in four Illinois harbors, 1999-2014.

List of Figures

Figure 1. Locations of Lake Michigan harbors sampled in 2014.

Figure 2. Mean lengths (± 1 S.E.) of Stock size (≥ 180 mm) smallmouth bass sampled under two different angling regulations: catch and release only (1998–2003, crosshatched) and a minimum size limit of 21 inches (2004–present, solid color).

Figure 3. Length distribution of Stock size (≥ 180 mm) smallmouth bass sampled at four Illinois harbors in 2014.

Figure 4. Mean lengths (± 1 S.E.) of Stock size (≥ 200 mm) largemouth bass sampled under two different angling regulations: catch and release only (1998–2003, crosshatched) and a minimum size limit of 21 inches (2004–present, solid color).

Figure 5. Length distribution of Stock size (≥ 200 mm) largemouth bass sampled at four Illinois harbors in 2014. No largemouth bass were sampled at Calumet Harbor.

Executive Summary

Electrofishing methods were used to assess the nearshore fish community in three Illinois harbors and the shoreline inside Calumet Harbor in summer 2014. North Point Marina had the highest number of species ($N = 17$), followed by Waukegan South Harbor ($N = 13$). Twelve species were found in Jackson Harbor and only eight in Calumet Harbor. Rock bass, largemouth bass, bluegill, pumpkinseed, and smallmouth bass were the most abundant target species that we encountered. Several of these species have expanded their ranges and are more actively pursued by anglers compared to a decade ago. For example, young-of-the-year smallmouth and largemouth bass have been collected recently by seining in areas at which no black bass have been stocked or juvenile bass have been collected in the past. In 2014, 127 smallmouth bass were of Stock size (≥ 180 mm), 85 of Quality size (≥ 280 mm), 56 of Preferred size (≥ 350 mm), 26 of Memorable size (≥ 430 mm), and no Trophy-size (≥ 510 mm) were sampled. Largemouth bass were similarly abundant but generally smaller than smallmouth bass. We sampled 189 largemouth bass of Stock size (≥ 200 mm), 117 of Quality size (≥ 300 mm), 50 of Preferred size (≥ 380 mm); no Memorable (≥ 510 mm) or Trophy (≥ 630 mm) sized largemouth bass were sampled. A black bass catch-and-release regulation for Lake Michigan, adopted in 1998, and subsequently changed to one black bass over 21 inches, has not resulted in large increases in catch-per-unit-effort or mean lengths of largemouth bass or smallmouth bass. Only one PIT-tagged black bass was collected in 2014; this largemouth bass was originally tagged in 2006 and grew approximately 4 inches in length over the eight years at liberty.

Introduction

Several sport and non-sport fish species inhabit Illinois harbors and nearshore areas of Lake Michigan in summer. Sport fish species found in these areas include: smallmouth bass (*Micropterus dolomieu*), largemouth bass (*Micropterus salmoides*), yellow perch (*Perca flavescens*), northern pike (*Esox lucius*), rock bass (*Ambloplites rupestris*), and several other Centrarchids (sunfish family). There has been an increasing demand for sport fishing opportunities in nearshore areas and an increased interest in the nearshore sport fishery since 1998, especially for black bass. Increases in the abundances of these warm- and cool-water fish species and angler effort for non-perch and non-salmonid fish species in the Illinois waters of Lake Michigan are evident from sport angler creel data. So few smallmouth bass were found in the possession of anglers that no estimate of harvest could be calculated from creel data prior to 1996; in 2000 however, anglers reported catching an estimated 4,892 smallmouth bass (pers. comm. W. Brofka, INHS).

Although management of fish species inhabiting these nearshore areas has been incorporated into the Illinois Strategic Plan for Lake Michigan fisheries since the early 1980s, personnel and funding deficiencies did not allow their investigation until 1995. This assessment program was developed to monitor the relative abundance and distribution of nearshore sport fish species and to determine whether those species were susceptible to overexploitation by tracking changes in relative abundances over time. Species composition, abundance, and length distribution data previously was obtained through incidental catches of non-salmonid fish species when sampling returning salmonids by electrofishing in the fall and through a sport angler creel survey. Sport anglers were interviewed, fish in their possession were measured for length and weight, and estimates of the sport harvest were considered to be estimates of the relative abundance of these fish species. Abundance and species composition data obtained through a creel survey, however, may be biased because anglers target specific species, effort is not equivalent at all locations, and harvest (rather than total catch) is usually reported. In addition to biological information (e.g., length and weight), an understanding of seasonal dispersal patterns of the sport fish associated with the nearshore fish community is required to effectively manage these species. If sport fish dispersal patterns for Lake Michigan are similar to the patterns

observed in Lake Ontario, then some of these fish species will inhabit protected areas early in the year and later move into open lake areas once water temperatures reach 15° C (Danehy 1984).

The objectives of this ongoing study are to: 1) determine the fish species composition of select Illinois harbors and nearshore areas of Lake Michigan; 2) monitor changes in the relative abundances of smallmouth and largemouth bass and other sport fish through time; 3) evaluate intra- and inter-annual fidelity of smallmouth and largemouth bass to harbors; 4) monitor growth indices for sport fish inhabiting these harbors; and 5) collect age-composition data which may eventually be used to determine recruitment rates of the most abundant fish species.

Methods

Fish were sampled using a Smith-Root Model 5.0 GPP Pulsed-DC electrofishing unit, operated at 8-10 amps and 60 pulses per second. Total sampling time was based on harbor size, weather conditions, and the amount and type of fish collected. Selection of sampling sites (Figure 1) was based on harbor configurations that were conducive to electrofishing (i.e., availability of shallow water areas <3 m in depth), availability of a launch ramp, and sport-angler creel survey data. Three Illinois harbors and the shoreline inside Calumet Harbor were selected for sampling in 2014 (Table 1). Sampling at North Point Marina was limited to the inner entrance of the north harbor, the channel connecting the north and south harbors, and the south harbor. At Waukegan, the south harbor was sampled, and the inner harbor was sampled at Jackson Harbor. The Calumet Harbor site consisted of the rip-rap shoreline between the Calumet River and the north slip within Calumet Harbor. The south face of the Calumet breakwall and the harbor at Calumet Harbor's Yacht Club were sampled on a few occasions.

Sport fish species were the target of electrofishing sampling effort. We attempted to capture all largemouth and smallmouth bass that were encountered; other target species were subsampled to obtain a representative distribution of sizes. Abundance of non-target species (e.g., alewife, gizzard shad, white sucker, and carp) was usually only noted. Sampled fish were dip-netted and held onboard in a 100 ga tank filled with a 0.5% solution of NaCl and lake water. An oxygen cylinder with an air stone was used to increase retention time and keep the fish alive

until biological data were obtained. Fish were measured to the nearest 5 mm (maximum total length) and weighed to the nearest 10 grams.

Relative Stock Density (RSD) for Quality, Preferred, and Memorable length fish were calculated for smallmouth and largemouth bass (Table 2; Gabelhouse 1983 as reported in Anderson and Gutreuter 1996).

Results and Discussion

North Point Marina had the highest number of species ($N = 17$), followed by Waukegan South Harbor ($N = 13$). Twelve species were found in Jackson Harbor and eight were sampled in Calumet Harbor. Rock bass, largemouth bass, bluegill, pumpkinseed, and smallmouth bass were the most abundant target species that we encountered. Overall, we sampled 12 sport (target) fish species by electrofishing in 2014, in addition to eleven non-target species (Table 3). It is likely that increased water clarity and aquatic vegetation in the protected areas of these harbors have produced favorable conditions for a number of these cool- and warm-water fish species (Jude et al. 2002). While Calumet still had a number of target species, very few Centrarchid species (the exception being rock bass) were present. This is likely a result of the Calumet site being an open-lake area and more exposed to wave action and rapid changes in water temperatures, which are not conducive for establishing aquatic vegetation.

The types of sport fish species we encountered in the three protected harbors were similar to those that are typically found in warm-water inland lakes with similar habitats. One major difference between these harbors and inland lakes is the abbreviated growing season in the harbors caused by influxes of cool water from the main lake, which suppresses water temperatures in the spring and may intermittently decrease temperatures during the summer due to upwellings. A second difference is the relatively large areas of restricted fishing access; much of the harbors' areas contain moored vessels and are closed to fishing. Thus, Illinois harbors may act as refuges on Lake Michigan where populations of warm-water fish may grow in a near natural state with limited fishing mortality.

Largemouth bass, smallmouth bass, bluegill, pumpkinseed, and rock bass were generally the most abundant target species that we encountered. Furthermore, some fish species were found only in particular harbors. For example, warmouth were collected only in North Point Marina and black crappie and northern pike were collected only in North Point Marina and Waukegan South Harbor.

Smallmouth bass

Sporadic and limited stockings of juvenile smallmouth bass occurred in 1969 through 1984 at Diversey, Belmont, and Great Lakes Harbors (Table 4). Jackson Harbor was stocked in 1985, although young-of-the-year (YOY) smallmouth bass had been collected during annual beach seine sampling at that site since 1982. The harbor at Great Lakes Naval Training Center in North Chicago was stocked with YOY smallmouth bass in 1973, 1977, and 1984, but no YOY smallmouth bass were captured in that harbor from 1982-1986. YOY smallmouth bass however were collected at Great Lakes Harbor from 1987 until 1994, when the site was no longer sampled. Additionally, YOY smallmouth bass have been collected at Calumet (since 1991) and Waukegan (1997) where no stocking occurred. Thus, it is likely that smallmouth bass stocking either contributed to the establishment of sustainable populations or only marginally supplemented an existing population in the case of Jackson Harbor. Since YOY smallmouth bass have been collected recently in areas at which no smallmouth bass have been collected in the past (e.g., Farwell Avenue Pier in 2000), it appears that the population also has expanded its range along the Illinois shoreline. A potential limiting factor to the further expansion of the smallmouth bass population may be insufficient spawning habitat (protected areas with stable, warm water), though relatively few successful nests can sustain a population (Coble 1975). Small Centrarchids (e.g., bluegill, pumpkinseeds) are known nest predators, but the presence of round gobies (*Neogobius melanostomus*) poses an additional threat to successful recruitment of smallmouth bass in Lake Michigan. The increased energy needed to successfully defend nests from predatory attacks by round gobies could deplete energy reserves quickly, potentially forcing smallmouth bass to abandon their broods (Steinhart et al. 2004).

We sampled 127 Stock size smallmouth bass in 2014 and 14 individuals less than Stock size (< 180 mm). Most smallmouth bass were collected at Calumet Harbor (N = 75). Similar

numbers of smallmouth bass were caught at North Point Marina (N=20) and Waukegan South Harbor (N=20), and only twelve were caught at Jackson Inner Harbor. Catch-per-unit-effort (CPE) declined at Waukegan South Harbor compared to 2013. CPE at the other three harbors increased although CPEs were below long-term averages (Table 5). Until 2007, a large number of bass tournaments held weigh-ins at Calumet Harbor where smallmouth bass were released after the weigh-in, therefore potentially inflating CPEs for sampling efforts; a decline in CPEs from 2007 onward may be attributed to the subsequent reduction of these tournament releases.

Mean length of smallmouth bass that were Stock-size (180 mm) or longer has been variable; mean length increased between 2008 and 2009, decreased in 2010 and 2011, rebounded in 2012 and 2013, and declined again in 2014 (Figure 2). Many smallmouth bass sampled in 2014 were of Quality size (N = 85; PSD = 67), fewer of Preferred (N = 56; RSD₃₅₀ = 44) size, and still fewer of Memorable (N = 26; RSD₄₃₀ = 20) size. No trophy-size smallmouth bass (\geq 510 mm) were sampled in 2014. Most young (< 250 mm) smallmouth bass were sampled at North Point Marina and Jackson Harbor; larger fish (>330 mm) were sampled at Calumet Harbor (Figure 3). There has been no distinguishable increase in mean length following the 1998 enactment of a “catch-and-release only” regulation for black bass in Lake Michigan and the change to “one fish \geq 21 inches” in 2004 (Linear Regression; $F_{1, 15} = 4.44$; $P > 0.05$).

Largemouth bass

No stocking records exist for largemouth bass, but nonetheless they are observed in the nearshore fish community. Populations may have occurred naturally in protected areas along the shoreline and dispersed from these remnant stocks. There are several potential sources for brood fish to have entered Lake Michigan in the past, such as the Lake Calumet complex, Wolf Lake, the Japanese Gardens ponds at 59th Street Harbor, Lincoln Park Zoo ponds, the diversion structure at the North Branch of the Chicago River (Wilmette), and Prairie Cove Harbor on the Illinois/Wisconsin state line. Remnant populations of brood fish may have existed in these locations until recent changes in the lake favored their dispersal.

Largemouth bass have been less actively pursued by anglers than smallmouth bass, yet Stock size largemouth bass (N = 189) were captured in greater numbers than smallmouth bass (N

= 127). CPEs in 2014 declined relative to 2013 at North Point Marina and Jackson Harbor (Table 6); no largemouth bass have been sampled at Calumet Harbor since 2009.

One PIT tagged largemouth bass was recaptured in 2014. This largemouth bass was tagged at North Point Marina in 2006 and was recaptured in at the same location. The fish grew from 275 mm (age-5) at tagging to 375 mm at recapture (age-13). Results of our monitoring efforts over the past 10 years indicate that there is high site fidelity to harbors. Like smallmouth bass, largemouth bass have been shown to move in and out of Illinois harbors, but typically not between them (Savitz and Treat 2007).

Many largemouth bass sampled in 2014 were of Quality size (N = 117; PSD = 62) and over one-quarter were of Preferred size (N = 50; RSD₃₈₀ = 26); no Memorable (510–625 mm) or Trophy (≥ 630 mm) largemouth bass were sampled (Figure 6). We sampled 91 individuals less than the Stock size (< 200 mm) in 2014. Between 2000 and 2014, mean lengths of largemouth bass significantly increased (Linear Regression; $F_{1,15} = 18.3$; $P < 0.001$). The most obvious distinction from visual inspection of the data was a sustained increase in mean length after 2006 (Figure 5). The data do not support definitive conclusions as to the specific reasons for this increase, but we speculate that a shift in prey consumption (e.g., addition of round goby as a large contributor to diets) or decreased competition for food may be responsible for the increase in sizes rather than the harvest regulation. Given low densities and a highly restrictive harvest regulations however, we would expect to sample larger fish in our surveys.

Recommendations

1. Monitor angler effort directed at smallmouth and largemouth bass and potential population expansions using shoreline creel surveys.
2. Collect a representative sample of abundant sport fish species biennially to determine ages using otoliths.

Acknowledgements

This study was conducted using Federal Aid in Sport Fish Restoration funds (grant number F-65-R).

Literature Cited

- Anderson, R. O., and S. J. Gutreuter. 1996. Length, weight, and structural indices. Pages 442–466 in Nielson, L. A., and D. L. Johnson (eds.) *Fisheries Techniques*. American Fisheries Society, Bethesda, Maryland.
- Brofka, W. A., and J. M. Dettmers. 2001. A Survey of Sport Fishing in the Illinois Portion of Lake Michigan. Aquatic Ecology Technical Report 01/2. 58 pp.
- Brofka, W. A., and J. M. Dettmers. 1999. A Survey of Sport Fishing in the Illinois Portion of Lake Michigan. Aquatic Ecology Technical Report 99/3. 57 pp.
- Brofka, W. A., and J. E. Marsden. 1996. A Survey of Sport Fishing in the Illinois Portion of Lake Michigan. Aquatic Ecology Technical Report 96/3. 52 pp.
- Coble, D. W. 1975. Smallmouth bass. Pages 21-30 in *Black Bass Biology and Management*.
- Cope, J.M. and A.E. Punt. 2007. Admitting ageing error when fitting growth curves: an example using the von Bertalanffy growth function with random effects. *Canadian Journal of Fisheries and Aquatic Sciences* 64: 205–218.
- Danehy, R. J. 1984. Comparative ecology of fishes associated with natural cobble shoals and sand substrates in Mexico Bay, Lake Ontario. MS Thesis. State University of New York, Syracuse. 84 pp.
- Fraser, J. M. 1955. The smallmouth bass fishery of South Bay, Lake Huron. *Journal of the Fisheries Research Board of Canada* 12:147–177.
- Hair, D. E. 1979. Some Aspects of the Life History of the Smallmouth Bass (*Micropterus dolomieu dolomieu*) in the Bass Islands Area of Lake Erie. Master's Thesis. Ohio State University. 57 pp.
- Jude, D., Stoermer, E., Johengen, T., and A. N. Perakis. 2002. Non-indigenous species in the Great Lakes: ecology, interactions, impacts, and future research directions. White paper prepared for the University of Michigan's Great Lakes Initiative. 39 pp.
- Savitz, J. and L. Treat. 2007. Movements and site fidelity of black bass in three harbors along the Illinois shoreline of Lake Michigan. *Journal of Freshwater Ecology* 22: 267–269.
- Steinhart, G. B., Sandrene, M. E., Weaver, S., Stein, R. A., and E. A. Marschall. 2004. Increased parental care cost for nest-guarding fish in a lake with hyperabundant nest predators. *Behavioral Ecology* 16: 427–434.

Table 1. Amount of electrofishing effort (hrs:min) and water temperatures in four Illinois harbor areas sampled in 2014. Only one harbor was sampled in May due to high winds, other field work, and a leak in the generator fuel tank.

	Location			
	North Point Marina	Waukegan South Harbor	Jackson Inner Harbor	Calumet Harbor
7 May 2014	na	na	0:42 / 56F	na
2 June 2014	0:55 / 64F	0:45 / 64F	na	na
9, 12 June 2014	1:00 / 62F	0:25 / 60F	0:34 / 68F	0:56 / 64F
17, 18 June 2014	0:55 / 62F	0:45 / 64F	0:40 / 72F	1:00 / 68F
2, 3 July 2014	0:55 / 60F	0:35 / na	0:32 / 80F	0:55 / 64F
21, 25 July 2014	1:05 / 70F	0:21 / 70F	0:36 / 73F	0:45 / 64F
7, 8 August 2014	0:55 / 74F	0:32 / 72F	na	0:22 / 71F

Table 2. Proposed minimum lengths (mm) for smallmouth and largemouth bass of various size categories (from Gabelhouse 1983).

Species	Size designation				
	Stock	Quality	Preferred	Memorable	Trophy
smallmouth bass	180	280	350	430	510
largemouth bass	200	300	380	510	630

Table 3. Fish species sampled during summer 2014 by electrofishing in three Illinois harbors and along the shoreline in Calumet Harbor. A = abundant or frequently sampled; P = few specimens observed or infrequently sampled.

	Location			
	North Point Marina	Waukegan South Harbor	Jackson Inner Harbor	Calumet Harbor
<i>target species</i>				
black bullhead	P	P	A	
black crappie	P	P		
brown bullhead			A	
bluegill	A	P	A	
green sunfish	P	P		
largemouth bass	A	A	A	
northern pike	P	P		
pumpkinseed	A	A	A	P
rock bass	A	A	A	A
smallmouth bass	A	A	P	A
warmouth	P			
yellow perch	P	P		
<i>non-target species</i>				
bluntnosed minnow	P		P	
brown trout	P	P		P
common carp			P	
coho salmon			P	
fat head minnow	P			
freshwater drum				P
golden redhorse				P
golden shiner		P	P	
rainbow trout	P		P	P
redhorse, misc.	P			P
white sucker	P	P		

Table 4. Smallmouth bass stocking in Lake Michigan harbors, 1969–1985.

Year	Location	Number
1969	Diversey Harbor	4,000
1969	Belmont Harbor	1,000
1969	Belmont Harbor	306
1971	Belmont Harbor, Diversey Harbor	5,000
1973	Great Lakes NTC Harbor	4,000
1977	Belmont Harbor	1,000
1977	Great Lakes NTC Harbor	5,000
1980	Diversey Harbor	20,000
1983	Lincoln Park Lagoon	25,000
1984	Great Lakes NTC Harbor	18,000
1985	Jackson Harbor	18,400

Table 5. Catch-per-unit-effort (No. fish / 1 hr electrofishing) of Stock-size (≥ 180 mm) smallmouth bass in four Illinois harbors, 1999-2014.

Year	Location			
	North Point Marina	Waukegan South Harbor	Jackson Inner Harbor	Calumet Harbor
1999	24.00	0.00	13.68	56.57
2000	27.60	0.80	6.00	26.50
2001	19.64	6.22	9.33	20.93
2002	16.81	6.91	10.67	42.67
2003	10.19	3.38	5.09	12.91
2004	13.67	1.64	7.57	57.04
2005	17.50	3.50	3.00	33.53
2006	10.89	9.66	1.33	34.83
2007	4.00	2.53	0.00	33.45
2008	8.83	8.67	2.78	30.29
2009	7.68	2.14	2.11	20.25
2010	5.92	0.45	2.40	25.00
2011	6.49	30.09	2.70	14.50
2012	5.77	7.12	1.46	20.27
2013	13.20	1.15	1.09	15.63
2014	5.91	3.48	3.91	18.91

Table 6. Catch-per-unit-effort (No. fish / 1 hr electrofishing) of Stock-size (≥ 200 mm) largemouth bass in four Illinois harbors, 1999-2014.

Year	Location			
	North Point Marina	Waukegan South Harbor	Jackson Inner Harbor	Calumet Harbor
1999	21.00	1.00	9.40	1.14
2000	32.00	17.20	30.67	0.00
2001	22.70	31.56	38.67	0.39
2002	22.16	23.63	42.67	0.67
2003	22.04	11.81	27.64	0.60
2004	47.33	37.16	68.11	0.74
2005	72.00	41.96	99.00	0.60
2006	35.78	29.41	53.33	1.12
2007	32.24	26.27	22.57	0.00
2008	50.53	14.67	49.07	0.96
2009	33.99	16.43	24.51	0.42
2010	24.87	8.51	26.40	0.00
2011	19.46	9.73	44.49	0.00
2012	10.99	7.12	19.02	0.00
2013	39.60	5.42	20.45	0.00
2014	32.81	6.23	16.96	0.00




Figure 1. Locations of Lake Michigan harbors sampled in 2014.




Figure 2. Mean lengths (± 1 S.E.) of Stock size (≥ 180 mm) smallmouth bass sampled under two different angling regulations: catch and release only (1998–2003, crosshatched) and a minimum size limit of 21 inches (2004–present, solid color).




Figure 3. Length distribution of Stock size (≥ 180 mm) smallmouth bass sampled at four Illinois harbors in 2014.




Figure 4. Mean lengths (± 1 S.E.) of Stock size (≥ 200 mm) largemouth bass sampled under two different angling regulations: catch and release only (1998–2003, crosshatched) and a minimum size limit of 21 inches (2004–present, solid color).




Figure 5. Length distribution of Stock size (≥ 200 mm) largemouth bass sampled at four Illinois harbors in 2014. No largemouth bass were sampled at Calumet Harbor.